



## Effect of Axial Angulation of Sacral Vestibule S1 on Morphometric Analysis of Sacral Vestibule Using Plain Computed Tomography in North West Indian Population

Tarun Kumar<sup>1</sup>, Narinder Singh<sup>2\*</sup>, Bhanu Awasthi<sup>3</sup>, Sunil K Raina<sup>4</sup>, Narvir S Chauhan<sup>5</sup>

<sup>1, 2, 3, 4, 5</sup> Dr. Rajendra Prasad, Government Medical College, Kangra at Tanda, Himachal Pradesh, India

DOI: <https://doi.org/10.33545/26648989.2019.v1.i1a.6>

### Abstract

**Background:** The present study was aimed to study and develop in-depth understanding of the effect of the axial angulation of sacral vestibule S1 on morphometry of sacral vestibule in North-West Indian population presenting to our institution, which will go a long way in planning to treat the posterior pelvic injuries with percutaneous screws, thereby reducing the morbidity associated with open fixation.

**Methods:** This study was conducted in the Department of Orthopaedics and Radiodiagnosis at Dr Rajendra Prasad Govt. Medical College, Kangra at Tanda over a period of one year. All the patients of the age >18 years and above submitting for either abdominal, lower spinal or non-orthopaedic pathology of pelvic region, presenting for computed tomography to the Department of Radiodiagnosis were included in the study. There was significant difference in axial angulation of S1 (P=0.000) and S2 (P=0.045) between males and females.

**Results:** The axial angulation of S1 ranged from 2° to 23.0° with a mean value of 11.01°±3.74°. There was a significant difference in coronal angulation of S1 between age-groups 18-30 and 51-60 years (10.43±3.10 vs. 11.26±3.74; P=0.023). There was a weak relation between axial angulation of S1 and age-groups 18-30 years (r=0.195; P=0.013), 31-40 years (r=0.139; P=0.201), 41-50 years (r=-0.103; P=0.237), 51-60 years (r=0.013; P=0.850), and >60 years (r=-0.150; P=0.642). There was a weak relation between interspinous distance with axial angulation of S1 (r=0.049; P=0.225). There was a weak relation between height with axial angulation of S1 (r=0.037; P=0.364).

**Conclusion:** The present study, the first of its kind in North Western part of India arrived to help us anthropometric measurements of sacral vestibule, thereby, helping in development of local protocols for percutaneous fixation in sacral fracture.

**Keywords:** sacral vestibule, Axial angulation of the vestibule, interspinous distance.

### Introduction

The sacral bone is an inverted triangle that sits obliquely between the two innominate bones of the pelvis at the distal aspect of the spinal column. It functions mechanically to convey axial load from the lumbar spine into the lower extremities for balanced locomotion. The ventral sacral body is concave and derived from five vertebrae. The transverse processes of the sacral vertebrae coalesce to form the sacral ala, which projects laterally from the upper sacral promontory [1].

The standard treatment of unstable sacral fractures is surgical fixation due to a high incidence of residual morbidity under conservative treatment. The primary goal is anatomic reduction, followed by a rigid fracture fixation. There are several operating techniques like fixation with iliosacral screws or plates, triangular osteosynthesis, ilioliac (plates, internal fixators, and bars) and trans-sacral screws or bars. In recent years, sacroiliac screws and spinopelvic internal fixators have become the preferred implants for fixation of posterior pelvic ring fractures. Whereas full weight bearing is allowed for most spinopelvic fixations, none or partial weight bearing is recommended for iliosacral screw fixations [2].

The present study was aimed to study and develop in-depth understanding of the morphometry of sacral vestibule in North-West Indian population presenting to our institution, which will go a long way in planning to treat the posterior

pelvic injuries with percutaneous screws, thereby reducing the morbidity associated with open fixation.

### Materials and Methods

This study was conducted in the Department of Orthopaedics and Radiodiagnosis at Dr Rajendra Prasad Govt. Medical College, Kangra at Tanda over a period of one year. All the patients of the age >18 years and above submitting for either abdominal, lower spinal or non-orthopaedic pathology of pelvic region, presenting for computed tomography to the Department of Radiodiagnosis were included in the study. The patients were informed about the aims and methods of the study and once consent was given for participation; they were evaluated. The evaluation included clinical assessment for height. This helped to draw comparison while arriving at morphometry of Sacral vestibule.

### The following patients were excluded from the study

1. Age < 18 years.
2. The patient with pelvic ring dysmorphism.
3. Osteolytic pelvic lesions.
4. Fractures involving the posterior elements.
5. Post operated cases of above fracture
6. Not willing to participate in the study
7. Implants obscuring the lumbosacral junction.

Each patient and his attendants were adequately informed about the aims, methods, the anticipated benefits and potential risks of the study and the discomfort it might entail them and the remedies thereof. Every precaution was taken to respect the privacy of the patient, the confidentiality of the patient's information and to minimize the impact of the study on his/her physical and mental integrity and personality. The patients were given the right to abstain from participation in the study or to withdraw consent to participate at any time of the study without reprisal. Due care and caution were taken at all stages of the research to ensure that the patient was put to minimum risk, suffer from no irreversible adverse effects and generally, benefit from and by the research. Written informed consent was obtained from all the patients and attendants included in the study.

The subjects included in this study followed the protocol generally used by the Department of Radiodiagnosis for the conditions mentioned above. The subjects were placed in the supine position with fully extended knee joint with patella facing the sky for CT examination. 3D volume reconstruction of surface anatomy of bony pelvis was then performed using available CT data.

All CT scan were included scanning by anteroposterior tomogram as well as axial images of Sacral Vestibule. All angles were measured at the CT workstation. Measurements of the angles were performed by a junior resident (the investigator) from the Department of Orthopaedics Dr. RPGMC Tanda and were supervised by consulting Orthopaedician and Radiologist.

Following parameters were noted in each patient according to the sex of the patient:


- Age of the patient
- Sex of the patient
- Axial angulation of vestibule S1
- Interspinus distance
- Height of the patient

**Statistical Analyses**

Data were presented as frequency, percentages, and median (inter quartile range; IQR). Difference between quantitative variables was compared using Mann Whitney U test. Spearman correlation coefficient was used to find relation between two variables. P value <0.05 was considered significant. Statistical analyses were performed using SPSS v20.

**Axial Angulation**

Axial angulation was measured as the angle subtended by a line drawn perpendicular to the axis of the osseous corridor and a line connecting the posterior iliac spines. The axis of axial CT reformats was reset so that the axis was perpendicular to the superior end plate of the first sacral segment. Reformats were then made perpendicular to the first sacral osseous corridor.



**Fig 1**

**Morphological Analysis of Sacral Vestibule Axial Angulation**

**Table 1**

	<b>S1</b>
Minimum	2°
Maximum	23°
Mean	11.01°±3.74°
Median	10
IQR	10

The axial angulation of S1 from 2° to 23.0° with a mean value of 11.01°±3.74°.

**Relation with age** (Comparative analysis of axial angulation of S1)

**Table 2**

	<b>Age-group</b>	<b>Mean±SD</b>	<b>Min-Max</b>	<b>P Value</b>
<b>S1</b>	18-30 <sup>a</sup>	10.43±3.10	-1-8	Pab=0.311; Pac=0.056
	31-40 <sup>b</sup>	10.87±3.51	-1-8	Pad=0.023; Pae=0.051
	41-50 <sup>c</sup>	11.28±4.41	-1-8	Pbc=0.475; Pbd=0.406
	51-60 <sup>d</sup>	11.26±3.74	-1-8	Pbe=0.199; Pcd=0.973
	>60 <sup>e</sup>	12.33±4.70	-1-8	Pce=0.430; Pde=0.341

There was a significant difference in coronal angulation of S1 between age-groups 18-30 and 51-60 years (10.43±3.10 vs. 11.26±3.74; P=0.023).

**Relation with Age**

**Table 2**

		<b>Correlation Coefficient (r)</b>	<b>P Value</b>
<b>S1</b>	18-30	0.195	0.013
	31-40	0.139	0.201
	41-50	-0.103	0.237
	51-60	0.013	0.850
	>60	0.150	0.642

There was a weak relation between axial angulation of S1 and age-groups 18-30 years (r=0.195; P=0.013), 31-40 years (r=0.139; P=0.201), 41-50 years (r=-0.103; P=0.237), 51-60 years (r=0.013; P=0.850), and >60 years (r=-0.150; P=0.642).

**Relation with Sex**

**Table 3**

		<b>Male</b>	<b>Female</b>	<b>P Value</b>
<b>S1</b>	Mean±SD	11.57±3.88	10.23±3.40	0
	Median	10	10	
	IQR	4	4	
	Median	3	4	
	IQR	5	5	

There was significant difference in axial angulation of S1 (P=0.000) between males and females.

### Relation with Interspinus distance

Table 4

	Correlation Coefficient (r)	P Value
S1	0.049	0.225

There was a weak relation between interspinus distance with axial angulation of S1 ( $r=0.049$ ;  $P=0.225$ )

### Relation with Height

Table 5

	Correlation Coefficient (r)	P Value
S1	0.037	0.364

There was a weak relation between height with axial angulation of S1 ( $r=0.037$ ;  $P=0.364$ )

### Discussion

The sacral vestibule concept has been proposed by Carlson *et al.* which is described as the three-dimensional screw space available in the narrowest part of iliosacral screw channel. The sacral vestibule is located in the transition zone between the sacral wing and sacral body and serves as the entrance to the sacral vertebrae [3]. The S1 vestibule, located above the sacral foramina and between the sacral foramina and the slope of the sacral wing, is the isthmus of the transition zone between the sacral wing and the S1 vertebrae. The present study was aimed to morphometrically analyze the axial angulation of sacral vestibule S1 using plain CT at Department of Orthopaedics, Dr RPGMC Kangra at Tanda. A total of 610 patients were included in the study.

The axial angulation of S1 ranged from  $2^{\circ}$  to  $23.0^{\circ}$  with a mean value of  $11.01^{\circ} \pm 3.74^{\circ}$ . There was a significant difference in coronal angulation of S1 between age-groups 18-30 and 51-60 years ( $10.43 \pm 3.10$  vs.  $11.26 \pm 3.74$ ;  $P=0.023$ ). There was a weak relation between axial angulation of S1 and age-groups 18-30 years ( $r=0.195$ ;  $P=0.013$ ), 31-40 years ( $r=0.139$ ;  $P=0.201$ ), 41-50 years ( $r=-0.103$ ;  $P=0.237$ ), 51-60 years ( $r=0.013$ ;  $P=0.850$ ), and  $>60$  years ( $r=-0.150$ );  $P=0.642$ ). There was a weak relation between interspinous distance with axial angulation of S1 ( $r=0.049$ ;  $P=0.225$ ). There was a weak relation between height with axial angulation of S1 ( $r=0.037$ ;  $P=0.364$ ).

The proper location and length, to insert iliosacral screws is parallel to the long diameter with the inclination angle of the vestibule; therefore, these both parameters are very important references for the operation. Due to smaller size in these parameters, the insertion location, direction of the screw, and the position relationships between the screws are particularly limited for female patients [4]. Sacral variations are common in Indian population; however, these parameters are higher than Chinese populations [5].

Kaiser *et al.* measurements showed that axial angulation of S1 vestibule was  $11 \pm 10.5^{\circ}$ . Our results are in concordance with Kaiser *et al.* [6]. Therefore, the placement of iliosacral screws should be considered carefully based on the size, gender, height and ethnicity of the patient.

### Conclusion

The present study, the first of its kind in North western part of India arrived to help us anthropometric measurements of

sacral vestibule, thereby, helping in development of local protocols for percutaneous fixation in sacral fracture.

### References

1. Sugar O. How the Sacrum Got Its Name. JAMA. 1987; 257:2061-3.
2. Mehta S, Auerbach JD, Born CT, Chin KR. Sacral fractures. J Amer Acad Orthop Surg. 2006; 14(12):656-65.
3. Carlson DA, Scheid DK, Maar DC, Baele JR, Kaehr DM. Safe placement of S1 and S2 iliosacral screws: the "vestibule" concept. J Orthop Trauma. 2000; 14(4):264-9.
4. Dong Q, Tian W, Ma B. Evaluation and clinical application of sacral S1 vestibule measurements in Chinese adults. Int. J Morphol. 2014; 32(1):202-7.
5. Singh ID, Ajita R, Singh N. Variation in the number of sacral pieces. IOSR Journal of Dental and Medical Sciences (IOSR-JDMS). 2015; 14:106-8.
6. Kaiser SP, Gardner MJ, Liu J, Routt ML Jr, Morshed S. Anatomic determinants of sacral dysmorphism and implications for safe iliosacral screw placement. J Bone Joint Surg Am. 2014; 96(e120):1-8.